Declassified in Part - Sanitized Copy Approved for Release 2012/11/01: CIA-RDP79B00873A000100010147-1

MEMORANDUM FOR:

IMAGE ANALYS PROGRAM

3 SEPARATE EXPERIMENTS

WITH EMULSION N.2.0056

(DATE)

Declassified in Part - Sanitized Copy Approved for Release 012/11/01 : CIA-RDP79B00873A000100010147-1

(47)

1 August 1968 25X1

∶25X1

Image Analysis Program TITLE:

Report No. TO-B 67-20 on Image Analysis

21 April 1967

Infectious Development Experiment: Consisting Of Counting The Silver Halide Crystals (Singles and In Clumps) In Exposed Undeveloped Sub-Mono Grain Layer Model Emulsion Number N-2-0056.

MORE SPECIFICALLY

WORK PROGRAM:

1. Photographed undeveloped film samples, from emulsion N-2-0056, under high magnification (1700x) and took a sufficiently large number of frames to encompass 9560 grains.

Micrographs were printed high contrast in the positive form on 8 x 10 inches D.W. Matt Photo Paper.

- 2. Determined how many single isolated grains there are and how many grains are situated inclumps by actually counting:
 - A. average total grains in all the eleven 8 x 10 photo prints (870 grains)
 - B. Average isolated single grains in all the eleven frames. (121 single grains)
 - C. Subtracted average total of single isolated grains from average total of all grains to yield total of grains in clumps. (749 in clumps)
 - D. Calculate average isolated singles as a percentage of average total grains counted in each 8 x 10 print.
 - 121 grains or 14% average isolated single grains counted in a given area. $(8" \times 10" \text{ Photo print at } 1700X)$
 - 3. See attachment 1.

1	Δ.,	***	1968
_	Au,	KUB U	7,400

25X1

WORK PROGRAM: Infectious Development Experiments Consisting Of Counting The Silver Halide Crystals (Singles and In Clumps) In Exposed Undeveloped Sub-Mono Grain Layer Model Emulsion Number N-2-0056.

Photomicrograph Plate Number	Silver Halide Isolated Single Crystals	Silver Halide `Isolated Single Crystals	Silver Halide Crystals in Clumps	Silver Halide Crystals in Clumps	
Number	Count	Per Cent	Count	Per Cent	
ı	112'	13%	729	87%	
1-A	130	16%	670	34%	
2	123	15%	712	85%	
3	120	14%	762	86%	
4	108	13%	750	87%	
5	120	13%	784	87%	
6	130	14%	763	86%	
7	134	15%	786	85%	
8	113	14%	712	86%	
9	110	13%	761	87%	
10	126	14%	805	86%	
Total 11 Plates Averages	1326 Singles 120.6	14%	8234 In Clum;	s 86%	

13 August 1968

25X1

25X1

TITLE: Image Ana

Image Analysis Program

Report No. TO-B 67-20 on Image

Analysis, Dated 21 April 1967

WORK PROGRAM: Infectious development experiment consisting of counting the

silver halide grains isolated singles and in clumps that were not developed in a 21 step density tablet made with sub-mono grain layer model emulsion No-2-0056 obtained

21 June 1968.


PROCEDURES:

- 1. This experiment consists of making seven photomicrographs of step 11 in a 21 step density tablet exposed on sub-mono'grain layer model emulsion, developed in D-19 5 minutes at 68° F, washed 31 minutes, treated to EK Co. Stop Bath 5 minutes, photo flo solution for 2 minutes and dried without swabbing. Step tablet not cleared in Hypo Fixing Bath.
- 2. Micrographs were printed high contrast in the positive form on 8×10 inches D.W. Matt Photo Paper.
- 3. Total grains undeveloped single isolated and those in clumps were counted.
- 4. Isolated undeveloped singles were counted and calculated to express percentage of total counted.
- 5. Any change in percent singles represents a development bias which arises from their difference in environment. The difference is a measure of the infectiousness of development.
 - 5. See attachments 1 and 2.

Declassified in Part - Sanitized Copy Approved for Release 2012/11/01 : CIA-RDP79B00873A000100010147-1

		CHMENT				•	•	ISA	uc.1968
	IMAG	E ANALY	SIS PROG	BRAM Marian	er englige	``	<u></u> .		25X1
	PLATE MERCENAN	Dews. Standard	TOTOLO LACADING			POTAL DEVELOR PRINTERS	SINGENTIACE /CE CANING SINGENTIACE /CE	SUNCENCED SUNCED SUNCED	1.61.02.22 00.72.00.00.00 00.73.00.00
Puor	12 12 None Control 12 12 12 12 12 12 12 12 12 12 12 12 12	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	Series Series			PACE OF STATE OF STAT	2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Cherrs Cours	10 10 10 10 10 10 10 10 10 10 10 10 10 1
1.	11.	.34	190	31	870	680	3.56%	i	į.
2.	11.	.34	189	34	870	681	3.90%	78.27%	
3,	11.	.34	196	28	870	674	3.21%	77.47%	
4.	11.	.34	177	26	870	693	2.98%	77.65%	
5.	_11,	.34	183	31	870	687	3.44 %	76.967	
٤.	11.	.341	214	47	870	656	5.40%	75.40Z	
7.	11.	.34	232	40	870	638	4.59%	73.03 %	

EMULSION #N·2·0056 D-19 5MIN. 68°F.



)	August	1968	
			25X1

25X1

TITLE: Image Analysis Program

Report # TO-B 67-20 on Image Analysis

dated 21 April 1967

WORK PROGRAM: Carry Out Infectious Development Experiments Consisting of Counting The Undeveloped Silver Halide Crystals Remaining In an Exposed, D-19 Developed, Unfixed, 21 Step Tablet of Sub-Mono Grain Layer Model Emulsion Number N-2-0056

PROCEDURE:

- 1. Make photomicrographs and print on 8×10 matt paper at 1700x most steps of a 21 step density tablet of sub-mono grain layer model emulsion number N-2-0056, that has been exposed to the light of a sensitometer, developed in D-19 for 5 minutes at $68^{\circ}F$, treated with Eastman Kodak Co. short stop solution, washed thoroughly in running water for 30 minutes, with 3 minutes immersion in E.K. Co. Photo Flo Solution and hung up to dry (without swabbing). This special film is not fixed in the usual hypo clearing solution.
- 2. Ignoring totally black grains, count total number of single (isolated) undeveloped grains and then the total number of undeveloped grains. Express the singles as a precentage of the total.
- 3. Plot percent singles as some function of exposure. Plot percent singles as a function of the percentage of all grains developed.
- 4. Any change in percent singles represent a development Bias which arises from their difference in environment. The difference is a measure of the infectiousness of development.

Work To Follow The Above Experiment:

Two additional step tablets were exposed identically to that mentioned above. One was developed in D-11 for 6 min. at 68°F and one was developed in D-8 for 3 min. at 68°F. Similar photomicrographs will be made of each of these step tablets and counted as outlined above to relate infectiousness to specific aspects or types of developer.

See Attachment 1 and 2.

Declassified in Part - Sanitized Copy Approved for Release 2012/11/01: CIA-RDP79B00873A000100010147-1

: -			d Сору Арр			2/11/01 : CIA		0873 <u>A</u> 00010	
	ice an	ALYSI	S EXPER	MENT		#N·2·005			25X
					. /4 8	6			\ 0.04.
	egi.	1/2 3		86 £ 100	1522		0	A ST X	
			Jan /		\$ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\				33
		Selection of the select	34 35°				THE SECOND STATES		3
7	STENERALINGERALI	DE. STEPHEN	Selection of the select	20 20 20 20 20 20 20 20 20 20 20 20 20 2		S. S	4 4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6		VEN SO DELL'ESTO
40	/	_	100	\ \\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	46643	1456	100,58	S/OFE	
	4	.27	560	80	870	310	9.19%	36%	
11-A	4	.27	5/2	84	870	348	7.65%	40%	100 110 110 110 110 110 110 110 110 110
/2	5	.28	381	75	870	479	8,64%	55%	- 249 - 549 - 747
12-A	5	.28	485	95	870	375	10.91%	43%	
/3	6	.27	484	77	870	386	8.85%	44%	W.
13-A	6	.29	453	୯୫	870	417	7.81%	48%	
14	7	.30	304	64	870	566	7.35%	65%	
14-A	. 7	.30	400	90	870	470	10.33%	5420	
15	8	.31	309	72	870	561	8.26%	61%	
15-A		.31	393	86	870	477	9.88%	55%	
16	9	.32	280	70	870	590	8.04%	68%	
16-A	9	.32	302	58	870	568	6.66%		993 39
17_	10	·33	244	40	870	626	4.59%	72%	
17-A	10	.33	244	44	870	626	5.05%	72%	• 1
13	11	.34	215	35	870	655	4.02%	. 1	•
19	12	.35	160	36	870	710	4.13%		
20	13	.36	131	21	870	739	241%		:
21	14	.37	138	16	870		1.83%		
22						732	1.26%		
ľ.	15	.38	101	.//	870	769	0.68%	ì	
23	16	.39	59	6. J	870	811	0.60%	13/0	

